




Hard Rock Water Jet Mining, a Novel Method to Extract Water from Poly-Hydrated Sulphates on Mars



Picture Credit: MTU/Honeybee
Robotics/Jetportal SFC

6/13/2018
SRR/PTMSS

Jeff Allen (PI), **Paul van Susante (Co-PI)**, Ezequiel Medici, Timothy Eisele, Kris Zacny (HBR), Julie Kleinhenz (NASA GRC)
+ many students




Michigan Tech

ESI17

Award: 80NSSC18K0252

HONEYBEE ROBOTICS
Spacecraft Mechanisms Corporation





ISRU Resource Trade Tree

Use ISRU to generate consumables / propellant (source: HLS²)

Equator

Mid Latitude (45 – 55 Deg)

Polar

Use Atmosphere

Use Mineral Deposits

D: 'Common' Regolith

C: Phyllosilicates

B: Poly-hydrated Sulfates

Use Water Ice Deposits

A: Buried Glacial Ice

~~Recurring Slope Lineae~~

Accessibility and minimal volume


~~Deep Groundwater~~

No Evidence of Existence


~~Permafrost~~

~~High Latitude Ice~~


*Outside of Acceptable
Human Landing Sites*



Key Characteristics by Feedstock




- Gypsum deposits would have the lowest mass AND power requirements of the granular deposits. Ice mining power not established due to less experience and available data.
- Typical martian regolith processed at low temperatures doesn't result in lower power (due to production rates) AND requires more mass -> NO ADVANTAGE



Gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) is promising

- Exists on Mars in all hydration forms (Gypsum, Basanite, Anhydrite)
 - Gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$)
 - Alpha and beta hemihydrate ($\text{CaSO}_4 \cdot 0.5\text{H}_2\text{O}$)
 - Anhydrite (CaSO_4)
- Exists in many places
 - Noctis Labyrinthus, Melas Chasma, Columbus Crater, Gale Crater, Mars poles etc.
- Promising source of water
 - 20.9% water by mass
 - Water bound in crystal, disassociates around 150 °C
 - Soft (Moh's Hardness of 2, 3-60MPa)



Mohs Hardness Scale			
	Mineral Name	Scale Number	Common Object
Increasing Hardness ↑	Diamond	10	
	Corundum	9	Masonry Drill Bit (8.5)
	Topaz	8	
	Quartz	7	Steel Nail (6.5)
	Orthoclase	6	
	Apatite	5	Knife/Glass Plate (5.5)
	Fluorite	4	
	Calcite	3	Copper Penny (3.5)
	Gypsum	2	Fingernail (2.5)
	Talc	1	






Surface occurrences of gypsum

(on Earth)





- Selenite tower gypsum crystal
- 4-5 inch tall



- Desert rose gypsum crystal
- 2 inch diameter



Other occurrences of gypsum



National geographic picture of Cave of Crystals in the Naica Mine in Mexico.

Gypsum curtains and stalagtites in Lechuguilla cave in New Mexico as part of the Carlsbad Caverns National Park



Michigan Tech

ESI17

Award: 80NSSC18K0252

HONEYBEE ROBOTICS 
Spacecraft Mechanisms Corporation



White Sands, NM (gypsum sand dunes)

- Gypsum dunes exist in the Martian Polar areas




Typical Excavation Methods

- Granular vs. Hard rock (ice cemented)



- Has to be already granular
- Large reaction forces
- Wear / maintenance
- Operation for long periods
- Needs crushing/milling







Classic Gypsum Mining Process (Earth)




Natural state




Drill blast holes




Load explosives




Blast



Load & Transport



Crush / Mill rocks





Calcine

Discard dehydrated gypsum






Size Reduction on Earth



- For drywall production needed
 - Particles <100 micron before Calcination






Michigan Tech

E



Alternate Excavation & Calcination Process



Advantages

No mechanical excavation and comminution = no wear


Use water for

Liberation,
Comminution
Transportation
Separation

To extract
MORE water

Early Stage Innovation Project


- Started Jan 15, 2018
- 3 years
- Understanding
 - Gypsum (contamination)
 - Sealing / Enclosure
 - Excavation
 - Particle sizes generated
 - Slurry Pumping
 - Particle separation
 - Water extraction
 - Thermal & Fluid flows





Gypsum





- Deposit Mechanism
 - Sabkha (tidal flats)
 - Deep basin
- Porosity/permeability
- Thermal properties
- Mechanical properties
- Variability





Sealing / Enclosure


- Exploring options
 - Double ring of 'saw edge'
 - Silicone fingers in between





Excavation

- 20 minute yield varied from a few oz. to 11 lbs (@ 2200 psi and 15 deg nozzle)








Particle Sizes Generated

- Effect of spray intensity
- 2700 psi
- 1.7 GPM
- Zero degree vs 15 degree





2700 psi, 1.7 GPM, 30 seconds											
Combined	25 degree		15 degree		zero degree						
	Green (25 degrees)		White (15 degrees)		Red (zero degree)						
Microns	percent of start	retained on sieve (g)	percent of start	retained on sieve (g)	percent of start	retained on sieve (g)					
2000.0	14.24	27.06	16.53	42.98	58.56	511.20					
1400.0	0.76	1.44	0.71	1.84	1.74	15.18					
1000.0	0.61	1.15	0.75	1.94	1.10	9.64					
850.0	0.33	0.62	0.42	1.08	0.52	4.52					
500.0	1.25	2.38	1.96	5.10	1.49	13.04					
250.0	3.58	6.80	7.32	19.04	4.01	35.00					
106.0	19.38	36.82	31.48	81.84	20.29	177.14					
0.0	3.67	6.98	7.95	20.66	0.74	6.46					
		84.84		176.3		772.9	total started				
		83.25		174.48		772.18	total recovered				
		190.00		260.00		873.00	mass on paper towel				
	56.18	106.75	32.89	85.52	11.55	100.82	left on paper towel (small particles)				





Particle Sizes Generated





Slurry Pumping



- Currently using Sump pump
- Particles settle quickly
- May require agitation using mixer or waterspray
- Sand particles found their perfect hole to get stuck in
- If using screens → need method to clean





Particle Separation


- 98% solid removal at >40 micron
- 4-10 GPM



Gypsum = 2.3 g/cc



Water Extraction



Product	Process	Options	Temp Start of process	Stages
Alpha Hemihydrate	Fully saturated environment	Steam Autoclave	43-49 C, above 97 C	110-130 C, end at 160 C
Beta hemihydrate	Under-saturated environment	Rotary kiln, kettle, flash calcining	43-49 C	116-121 C 1 st boil, 149-166 C Continuous up to 140-154 C Batch up to 150-165 C
Soluble anhydrite	Keep heating either one		177 C 2 nd boil	Up to 210 C
Dead burned anhydrite	Keep heating either one		210 C	Up to 540 C

Dry heating issue

110 g gypsum got
16 g water
(22 g max)

Pyrex (820 C)
softening point






Modeling: Analytical & Pore Network Model

Coarse pores: 400 pores




$T_0 = 20\text{ C}$

$P_0 = 2\text{-}3\text{ kPa}$

$T_{\text{inf}} = -55\text{ C}$




Frost line at $T_{\text{inf}} = -55^\circ\text{C}$






Conclusion

- No show stoppers found
- Large variability of natural rock material
- Component requirement definition ongoing
- Starting system analysis / synthesis
- Mining process for detailed conops
- Future expansion:
 - Combine with gas transport
 - Shopvac / dental setup
 - Works with granular, ice-cemented and hard rock





Available Facilities MTU – Looking for ISRU Partners

- Rock crushing lab
- Vacuum chambers
- Chemical Processing Operations lab
- Modeling capabilities (super computer)
- KRC
 - 900 acres test terrain (all year)
 - Cold, ice, snow, winter testing
 - Many different terrains, slopes, surface features
 - Multiple 18-wheeler bays climate controlled (-80C)
 - Mobility, robotics, detection etc.
 - Basalt quarry

April 2018





Acknowledgements

This work was supported by an Early Stage Innovations grant from NASA's Space Technology Research Grants Program

- Many students:
 - Shahab Bayani Ahanger (Grad Student)
 - Wonyoung Choi (Highschool Student)
 - Michael Foetish (Grad Student)
 - Preston Hogue (Undergrad Student)
 - Udit Sharma (Grad Student)
 - Andrew Watson (Undergrad Student)
 - Mining INnovation Enterprise students (12)

