

# Modeling The Thickness And Shape Of Lunar Volatile Stability Zones

---

Hunter Danque and Kevin Cannon

Thursday June 9, 2022

# Outline

- **Introduction**
- **Background**
- **Methods**
- **Results**
  - **Physical and Thermal models**
  - **Topographic diffusion and real crater profiles**
- **Conclusions**



# Hypotheses



- Volatiles may experience subsurface temperatures that could sublimate ice from the base of cold traps
- Temperature conditions may exist to redeposit and concentrate volatiles in predictable parts of subsurface cold traps
- There may be a positive feedback mechanism in which increased ice content at the base of volatile stability improves the thermal conductivity and thickens volatile stability zones

# Background – Cold Traps

- The water ice stability zone exists where:
  - regolith surface temperatures are below ~110 K
  - subsurface temperatures are below **~145 K**
  - these conditions occur in some polar regions on the Moon
- They are often found in Permanently Shadowed Regions (PSRs) that may host cold traps
- The threshold for defining a cold trap is a sublimation rate of  $1 \text{ kg m}^{-2} \text{ Ga}^{-1}$






# Illumination and Temperature of a PSR






# Background – Obliquity and TPW

- About  $3 \pm 1$  Ga the Moon was at half its current semimajor axis
- The Moon may have experienced a Cassini State 1 to 2 transition and experienced obliquity as high as 77 degrees and extended periods at 25-50 degrees
- The Moon may have experienced true polar wander (TPW) with evidence for a paleopole at about 3.5 Ga
- During TPW the planet/moon realigns the rotation axis to the maximum principal axis of inertia when there is a change in internal mass distribution




# Background – Hydrogen and Ice Depth


- Hydrogen in upper meter from epithermal neutron data (a. b.)
- Calculated current and paleo pole depth to ice stability (c. d.)
- Combined current and paleo pole ice depths match the hydrogen content well (e. f.)




# Background - Topographic Diffusion



- Example of craters change in infill thickness and slope as a function of time
- $\frac{dh}{dt} = \kappa \nabla^2 h$ , where  $\kappa$  is the diffusivity




- Conceptual setup of a small impactor on the slope of a larger crater or hillside
- There is a net translation of mass in the downslope direction





# Methods-Model Properties

- Thermal conductivity (k)
  - Ice @ 110 K =  $5.5 \text{ W m}^{-1} \text{ K}^{-1}$
  - Regolith =  $0.023 \text{ W m}^{-1} \text{ K}^{-1}$
  - Upper Megaregolith =  $0.2 \text{ W m}^{-1} \text{ K}^{-1}$
  - Lower Megaregolith =  $1.0 \text{ to } 2.0 \text{ W m}^{-1} \text{ K}^{-1}$
- Density
  - Ice @ 110 K =  $932 \text{ kg m}^{-3}$
  - Regolith =  $1100\text{-}1800 \text{ kg m}^{-3}$
  - Upper Megaregolith =  $1960\text{-}2600 \text{ kg m}^{-3}$
  - Lower Megaregolith =  $2600\text{-}2900 \text{ kg m}^{-3}$



# Initial Synthetic Crater

Thermal diffusivity

$$\alpha = k / \rho c_p$$

$k$  is thermal conductivity


$\rho$  is density


$c_p$  is heat capacity

Upper Boundary conditions are:

- 110 K cold trap
- 160 K illuminated

Lower Boundary condition is 90 K/km




 Original water ice stability zone

Int. Crater= -181 m

160 , Temperature (K)  
0.023 , Thermal Conductivity (W/m/K)  
1800 , Input Density (kg/m<sup>3</sup>)  
448 , Heat Capacity (J kg<sup>-1</sup> K<sup>-1</sup>)  
2.84e-08 , Thermal Diffusivity (m<sup>2</sup>/s)


110 , Temperature (K)  
0.023 , Thermal Conductivity (W/m/K)  
1800 , Input Density (kg/m<sup>3</sup>)  
307 , Heat Capacity (J kg<sup>-1</sup> K<sup>-1</sup>)  
4.15e-08 , Thermal Diffusivity (m<sup>2</sup>/s)

# Filled Synthetic Crater






# Filled Synthetic Crater





# Filled Synthetic Crater with ice






# Crater - VIPER Mission Area




Ejecta thickness  $t = 0.14 R^{0.74} (r/R)^{-3.0}$






# Crater - VIPER Mission Area



# Lunar volatile system timeline – VIPER area example



# Conclusion

- The results appear to support potential for remobilization and concentration of water ice near the base of volatile stability
- Future work will develop and improve:
  - Physical crater fill thickness estimates from diffusion and ejecta fill
  - Volatile concentration scenarios



# References

- Carnahan, E., Wolfenbarger, N.S., Jordan, J.S., Hesse, M.A., 2021. New insights into temperature-dependent ice properties and their effect on ice shell convection for icy ocean worlds. *Earth and Planetary Science Letters* 563, 116886. <https://doi.org/10.1016/j.epsl.2021.116886>
- Fassett, C.I., Beyer, Heldmann, L., Hirabayashi, T., Keszthelyi, L., Lim, D.S.S., 2022. Age Estimates for Permanently Shadowed Craters in the Viper Mission Area. Presented at the 53rd LPSC, p. 2.
- Fassett, C.I., Crowley, M.C., Leight, C., Dyar, M.D., Minton, D.A., Hirabayashi, M., Thomson, B.J., Watters, W.A., 2017. Evidence for rapid topographic evolution and crater degradation on Mercury from simple crater morphometry. *Geophysical Research Letters* 44, 5326–5335. <https://doi.org/10.1002/2017GL073769>
- Fassett, C.I., Thomson, B.J., 2014. Crater degradation on the lunar maria: Topographic diffusion and the rate of erosion on the Moon. *Journal of Geophysical Research: Planets* 119, 2255–2271. <https://doi.org/10.1002/2014JE004698>
- Heiken, G.H., Vaniman, D.T., French, B.M., 1991. *Lunar sourcebook - A user's guide to the Moon*. Cambridge University Press.
- Keihm, S.J., Langseth, M.G., 1975. Microwave Emission Spectrum of the Moon: Mean Global Heat Flow and Average Depth of the Regolith. *Science* 187, 64–66. <https://doi.org/10.1126/science.187.4171.64>
- McGetchin, T.R., Settle, M., Head, J.W., 1973. Radial thickness variation in impact crater ejecta: implications for lunar basin deposits. *Earth and Planetary Science Letters* 20, 226–236. [https://doi.org/10.1016/0012-821X\(73\)90162-3](https://doi.org/10.1016/0012-821X(73)90162-3)
- Rücker, C., Günther, T., Wagner, F.M., 2017. pyGIMLi: An open-source library for modelling and inversion in geophysics. *Computers & Geosciences* 109, 106–123. <https://doi.org/10.1016/j.cageo.2017.07.011>
- Schorghofer, N., Aharonson, O., 2014. The Lunar Thermal Ice Pump. *ApJ* 788, 169. <https://doi.org/10.1088/0004-637X/788/2/169>
- Siegler, M., Nagihara, S., Grott, M., Warren, P.H., Paige, D.A., Kiefer, W.S., Smrekar, S.E., Wiczorek, M.A., 2020. Lunar Heat Flow: Global Predictions and Reduced Heat Flux (Working Paper).
- Siegler, M.A., Bills, B.G., Paige, D.A., 2011. Effects of orbital evolution on lunar ice stability. *Journal of Geophysical Research: Planets* 116. <https://doi.org/10.1029/2010JE003652>
- Siegler, M.A., Miller, R.S., Keane, J.T., Laneuville, M., Paige, D.A., Matsuyama, I., Lawrence, D.J., Crotts, A., Poston, M.J., 2016. Lunar true polar wander inferred from polar hydrogen. *Nature* 531, 480–484. <https://doi.org/10.1038/nature17166>
- Soderblom, L.A., 1970. A model for small-impact erosion applied to the lunar surface. *Journal of Geophysical Research (1896-1977)* 75, 2655–2661. <https://doi.org/10.1029/JB075i014p02655>
- Ward, W.R., 1975. Past Orientation of the Lunar Spin Axis. *Science* 189, 377–379. <https://doi.org/10.1126/science.189.4200.377>
- Warren, P.H., Rasmussen, K.L., 1987. Megaregolith insulation, internal temperatures, and bulk uranium content of the moon. *Journal of Geophysical Research: Solid Earth* 92, 3453–3465. <https://doi.org/10.1029/JB092iB05p03453>
- Zhang, J.A., Paige, D.A., 2009. Cold-trapped organic compounds at the poles of the Moon and Mercury: Implications for origins. *Geophysical Research Letters* 36, 5.