

Retrieving Subsurface Properties of Mars-Analog Glaciers with Drone-Based GPR

Roberto J. Aguilar¹ (roberto@lpl.arizona.edu), Tyler M. Meng², Michael S. Christoffersen³, Stefano Nerozzi¹, Jack W. Holt^{1,4}



UA SCIENCE
**LUNAR & PLANETARY
LABORATORY**

¹Lunar and Planetary Laboratory, University of Arizona, ²Department of Earth, Environmental and Planetary Sciences, Washington University in St. Louis, ³School of Earth and Atmospheric Sciences, Georgia Institute of Technology, ⁴Department of Geosciences, University of Arizona.


1. Goals

- Investigate the detectability of supraglacial debris, glacial base, and internal layering in terrestrial debris-covered glaciers using a drone-based ground penetrating radar (DGPR)
- Compare the results of DGPR with surface-based GPR at multiple frequencies
- Validate the detection of subsurface reflectors with clutter simulations

2. Background

Mars


Earth




Debris-covered glacial landform DEM from HiRISE and CTX



Sourdough Rock Glacier, Alaska. Airborne photo by Eric Petersen




DCGs on Mars ^[1]



DCGs on Earth ^[2]
(excluding Antarctica)


- Lobate debris aprons (LDAs) are debris-covered glacial landforms at the mid-latitudes of Mars, formed in multiple episodes during the Amazonian age [0.3 – 1 Ga] ^[3,4]
- The radar sounder SHARAD (15–25 MHz) onboard the Mars Reconnaissance Orbiter (MRO) has confirmed that the bulk composition of LDAs is nearly water ice ^[5]
- SHARAD hasn't directly resolved the supraglacial debris to ice contact or near surface stratigraphy ^[6]
- On Earth, debris-covered glaciers (DCGs) are planetary analogs for LDAs



Model for the formation of Galena Creek Rock Glacier ^[7]

3. Platform

Drone-based ground penetrating radar in Galena Creek Rock Glacier



Platform:


- Drone DJI Matrice 600 Pro
- GPR MALA Geodrone 80
- Impulse radar
- Center frequency: 80 MHz
- Bandwidth: 40 MHz

Navigation:


- Automated flight path
 - Speed (1 m/s)
 - Altitude (3 m–5 m)
 - Real-time terrain following
- Total takeoff weight: 10 Kg**

4. Supraglacial debris and bulk thickness

Drone-based GPR (80 MHz)



Surface-based GPR (50 MHz)






Profiles at the toe of Sourdough Rock Glacier

We manually picked the following reflectors:

- Surface (first return)
- Debris/ice interface
 - Debris thickness calculated with a velocity of 0.1 m/ns
- Basal reflector
 - Bulk thickness calculated with a velocity of 0.15 m/ns

* Velocity was retrieved with a common-midpoint (CMP) survey ^[8]

5. Internal debris layering and clutter simulations



The clutter simulator ^[9-10] used as input the digital elevation models (DEMs) acquired with drone photogrammetry (5 cm/px) ^[11]

6. Conclusions

- We successfully employed a DGPR platform to survey terrestrial DCGs, resolving debris-ice contacts, internal glacier stratigraphy, and total glacial thickness.
- The DGPR platform and our results over Martian analog ISRU targets demonstrate the potential for drone-based planetary exploration.

7. Future work

Development of a drone-based chirped radar

- Test behaviour of current and future radars, with a frequency between 10 MHz to 1.1 GHz
- Flight altitude ~120 m, mass < 5 kg.
- within the specifications of the Mars Science Helicopter ^[12]

Acknowledgments: This work was funded by NASA Solar System Workings Grant 80NSSC19K0561 and by the France-Arizona Institute for Global Grand Challenges. We thank the members of the 2023 Resource Exploration and Science of OUR Cosmic Environment (RESOURCE) team for their contributions to data acquisition at Galena Creek Rock Glacier: Michael Daniel, Rocio Jacobo, Rishi Chandra, Daniel Thomas, Nathaniel Putzig, Aaron Russell, and Callum Walter.

References: [1] Hauber, E. et al. (2008) JGR Planets 113. [2] Scherler, D. et al. (2018) JRL. [3] Forget, F. et al. (2006) Science, vol. 311. [4] Fassett, C. I. et al. (2014) Geology, vol. 42. [5] Holt, J. W. et al. (2008) Science, vol. 322. [6] Baker, D. M. H. & Carter L. M. (2019) Icarus 319. [7] Petersen, E. I. et al. (2019) Journal of Glaciology, 66. [8] Meng, T. M., et al. (2023) Journal of Glaciology 69. [9] Choudhary P. et al. (2016) IEEE Geoscience and Remote Sensing Letters. [10] Christoffersen M. S. et al. (2024) github.com/lpl-tapir/simc. [11] Meng T. M. et al (2023) Remote Sensing 15, 4779. [12] Tzanetos, T. et al. (2022) IEEE Aerospace Conf., pp. 01-16.